

EPJ manuscript No.
(will be inserted by the editor)

The hypernuclear physics heritage of Dick Dalitz (1925-2006)

Avraham Gal

Racah institute of Physics, The Hebrew University, Jerusalem 91904, Israel

Received: date / Revised version: date

Abstract. The major contributions of Richard H. Dalitz to hypernuclear physics, since his first paper in 1955 to his last one in 2005 covering a span of 50 years during which he founded and led the theoretical study of hypernuclei, are reviewed from a personal perspective. Topical remarks on the search for quasi-bound \bar{K} -nuclear states are made.

PACS. 01.30.-y – 01.60.+q – 01.65.+g – 21.80.+a

1 Introduction

Dick Dalitz was born in Dimboola, in the state of Victoria, Australia, on February 28th 1925, and gained B.A. and B.Sc. degrees in Mathematics and Physics in 1944 and 1945, respectively, from the University of Melbourne. He moved to Britain in 1946 for postgraduate studies at Cambridge, and then worked at the University of Bristol before joining in 1949 Rudolf Peierls in Birmingham. There he completed and wrote up his Ph.D. thesis on ' $0^+ \rightarrow 0^+$ transitions in nuclei', supervised by Nicholas Kemmer of Cambridge, and subsequently became a Lecturer. He spent two years in the U.S. from 1953, holding research positions at Cornell and Stanford, visiting also Princeton and Brookhaven National Laboratory, and returned as a Reader in Mathematical Physics to the University of Birmingham for a year before becoming Professor of Physics in the Enrico Fermi Institute for Nuclear Studies and the Department of Physics at the University of Chicago in 1956. He moved to Oxford in 1963 as a Royal Society Research Professor, the post he held until his retirement in 1990. In addition to the Dalitz Plot, Dalitz Pair and the Castillejo-Dalitz-Dyson (CDD) Pole that bear his name, he pioneered the theoretical study of strange baryon resonances, of baryon spectroscopy in the quark model, and of hypernuclei, to all of which he made outstanding contributions. His formulation of the $\theta - \tau$ puzzle led to the discovery that parity is not a symmetry of the weak interactions. A complete bibliography of Dalitz's works is available in Ref. [1].

During his postgraduate studies he spent a year working alongside Cecil Powell's cosmic ray group at Bristol and it was during this period that he took particular interest in the strange particles that were beginning to appear in cosmic rays and at particle accelerators. These included the first hyperfragment in 1952 [2] which inspired a lifelong interest in hypernuclei. Later on, he made significant contributions to the strong interactions of the strange parti-

cles and their resonant states [3,4]. As early as 1959 Dalitz and Tuan, by analysing the data on the strong interactions of K^- mesons with protons, predicted the existence of an $I = 0$, $J^\pi = (1/2)^-$ strange resonance about 20 MeV below the K^-p threshold [5]. This $\Lambda(1405)$ resonance was discovered two years later in the Berkeley hydrogen bubble chamber, studying the reaction $K^-p \rightarrow \Sigma + 3\pi$ for several charge states [6]. The proximity of this s -wave $\pi\Sigma$ resonance to the $\bar{K}N$ threshold suggested that it can be generated by $\bar{K}N - \pi\Sigma$ inter-hadron forces, and this was shown in 1967 by Dalitz *et al.* to be possible within a dynamical model of SU(3)-octet vector-meson exchange [7] which is, in fact, the underlying physical mechanism for the Tomozawa-Weinberg leading term in the chiral expansion of the meson-baryon Lagrangian [8,9]. The vector mesons ρ, ω, K^*, ϕ , which were discovered in the years 1960-62, relying heavily on Dalitz plots for some of these, were unknown when the $\Lambda(1405)$ was predicted. In the years to follow, Dalitz repeatedly considered the completeness of this dynamical picture, whether or not the S -matrix pole of $\Lambda(1405)$ due to the inter-hadron forces need not be augmented by a CDD pole arising from inter-quark forces upon allowing for an intermediate uds configuration. It is here that the earlier CDD discussion [10] found a fertile physical ground.

Looking back years later at the development of his own career, he made the following remarks [11] (which he rarely allowed himself to make in public):

- *Yes, as Gell-Mann said, pion physics was indeed the central topic for theoretical physics in the mid 1950s, and that was what the young theoretician was expected to work on. The strange particles were considered generally to be an obscure and uncertain area of phenomena, as some kind of dirt effect which could not have much role to play in the nuclear forces, whose comprehension was considered to be the purpose of our research. Gell-Mann remarked that he spent the major*

part of his effort on pion physics in that period, and I did the same, although with much less success, of course.

- *Fashions have always been strong in theoretical physics, and that holds true today as much as ever. The young physicist who is not working on those problems considered central and promising at the time, is at a disadvantage when he seeks a post. This tendency stems from human nature, of course, but it is unfortunate, I think, that the system operates in such a way as to discourage the young physicist from following an independent line of thought.*

Although about 30% of his research papers were devoted or connected to hypernuclei, Dalitz was primarily a particle physicist. This is reflected in the interview he gave during HYP03 [12], where hypernuclei get only the following two brief remarks:

- *My interest in hypernuclear events developed particularly well in Chicago because a young emulsion experimenter, Riccardo Levi-Setti, whose work I had known from his hypernuclear studies in Milan, came to the Institute for Nuclear Studies at this time. We each benefited from the other, I think, and we got quite a lot done.*
- *I was responsible for organizing particle-physics theory in Oxford. Besides quark-model work, I still did work on hypernuclear physics, much of this with Avraham Gal of Jerusalem.*

I first met Dalitz as a young student attending the 1966 Varenna International School of Physics “Enrico Fermi”, Course XXXVIII on ‘Interaction of High-Energy Particles with Nuclei’. He gave a series of lectures on the status of Hypernuclear Physics, and I was lucky to have been able to intercept him during one of the lectures with an important omission he made in a calculation which my shell-model background education at the Weizmann Institute gave me an advantage over him in calculating it correctly. This was the beginning of a very close collaboration lasting about 20 years during which we would often meet for joint periods of work, always discussing the latest experimental results and their likely interpretations. I have been amazed at Dalitz’s encyclopaedic knowledge and mastery of measurements and calculations in particle physics and also of many aspects of nuclear physics, his critical assessment of experimental results and his thoroughness at work. He always insisted on and managed to calculate things in his own way, relying only on facts, never on fancy. Our ways parted off after 1985, but we still maintained a close relationship until very recently, when I edited his last publication, the talk he gave at HYP03 [13].

2 Λ hypernuclei

2.1 The beginning

Dalitz pioneered the theoretical study of hypernuclei. His first published work on Λ hypernuclei dates back to 1955,

titled *Charge independence in light hyperfragments* [14]. It focused on the near equality of the (${}^4_\Lambda\text{H}$, ${}^4_\Lambda\text{He}$) binding energies and its origin in the charge symmetry of the ΛN interaction, and on the exceedingly small binding energy of ${}^3_\Lambda\text{H}$, the only bound $A = 3$ hypernucleus marking the onset of Λ -hypernuclear binding. By 1959 his analyses of the light, s -shell hyperfragments led him to state [15] that *the existence of a bound Λ -nucleon system is strongly excluded* and that *the analysis of the $T = 1$ triplet ${}^3_\Lambda\text{He}$, ${}^3_\Lambda\text{H}$, ${}^3_\Lambda\text{n}$ indicates that these systems are not expected to form bound states*, and that these essential conclusions *would not be seriously affected if there exist moderately strong three-body forces arising from pion exchange processes*. He returned in 1972 to consider the possible effects of three-body ΛNN forces in the s shell [16] quantifying what has been since called ‘the overbinding problem’, namely that the binding energy of ${}^5_\Lambda\text{He}$ comes out too large by 2 – 3 MeV in any calculation that fits well the binding energies of the lighter hypernuclei.¹

In a series of works covering three decades, he used the main $\Lambda \rightarrow p\pi^-$ weak-decay mode of light hypernuclear species studied in emulsion and bubble chambers to determine their ground-state spins and, thereby, to gain information on the spin dependence of the ΛN force. When he had begun this line of works, just before parity violation was realised during the turbulent 1956-1957 period, he wrongly concluded in a talk given at the 6th Annual Rochester Conference on High Energy Nuclear Physics in April 1956 that the triplet ΛN s -wave interaction was stronger than the singlet one [17]. His argument was based on assuming that parity was respected in the weak decay ${}^4_\Lambda\text{H} \rightarrow \pi^- + {}^4\text{He}$. Since the final products all had spin zero, and the pion was known to have a negative intrinsic parity with respect to nucleons, (quoting Dalitz, in *italics*) *the spin-parity possibilities for the (${}^4_\Lambda\text{H}$, ${}^4_\Lambda\text{He}$) doublet are $0^-, 1^+, 2^-, \text{etc.}$* Assuming (at that time it was still uncertain) that the Λ hyperon had spin-parity $(1/2)^+$, the spin-parity of ${}^4_\Lambda\text{H}$ had to be 1^+ , and this meant that the triplet ΛN s -wave interaction was stronger than the singlet one, and *one also concludes that the spin-parity for ${}^3_\Lambda\text{H}$ is $(3/2)^+$* . Of course we now know that this was wrong; and indeed soon after Dalitz himself, realising the merits of the strong spin selectivity provided by parity violation in the weak-interaction pionic decays of Λ hypernuclei, calculated the branching ratios of the π^- two-body decays of ${}^4_\Lambda\text{H}$ and ${}^3_\Lambda\text{H}$ to the daughter ground states of ${}^4\text{He}$ and ${}^3\text{He}$, respectively, in order to determine unambiguously the ground-state spins of the parent hypernuclei [18] which in a few years became experimentally established as 0^+ [19] and $(1/2)^+$ [20] respectively. This led to the correct ordering of the triplet and singlet ΛN s -wave interactions as we understand it todate.

2.2 The later years

Dalitz’s work on the p -shell hypernuclei, dates back to 1963 when together with Levi Setti, in their only joint pa-

¹ this need not be the case once $\Lambda N - \Sigma N$ coupling is explicitly allowed in.

per [21], *Some possibilities for unusual light hypernuclei* were discussed, notably the neutron-rich isotopes of ${}^6_{\Lambda}\text{H}$ and ${}^8_{\Lambda}\text{He}$ belonging to $I = 3/2$ multiplets, but his systematic research of the p -shell hypernuclei started in 1967 together with me laying the foundations for a shell-model analysis of Λ hypernuclei. As early as 1969 data on excited states were reported with the Λ hyperon in a $(1p)_{\Lambda}$ state coupled to the nuclear ground-state configuration, first from emulsion data [22,23] observing proton decay in some special instances such as ${}^{12}_{\Lambda}\text{C}$, and later on through in-flight (K^- , π^-) experiments at CERN and BNL. In the particular case of the ${}^{12}_{\Lambda}\text{C}$ excited cluster of states about 11 MeV above the $(1s)_{\Lambda}$ ground state, Dalitz participated actively in the first round of theoretical analysis for both types of experiments [24,25]. However, confronting these and similar data posed two difficulties which we identified and discussed during 1976. The first one was connected to understanding the nature of the Λ continuum spectrum which, owing to the small momentum transfer in the forward-direction (K^- , π^-) reaction in flight, was thought to consist of well defined Λ -hypernuclear excitations. It was not immediately recognised that since the Λ hyperon did not have to obey the Pauli exclusion principle with nucleons, hypernuclear quasi-free excitation was possible even at extremely small values of the momentum transfer, a possibility that was pointed out and analysed quantitatively by us [26] following the first round of data taken by the Heidelberg-Saclay collaboration at the CERN-PS in 1975. The other difficulty was connected with understanding the role of coherent excitations in the $(1p)_{\Lambda}$ continuum, the so called ‘substitutional’ or ‘analogue’ states, where the early theoretical concept of analogue states stemmed from considerations of octet-SU(3) unitary symmetry. Already in his first discussion of these states in 1969 [27], Dalitz recognised *that the strong excitation of these states does not depend on SU(3) symmetry. In fact it is reasonable to believe that SU(3) symmetry has almost no relevance to the relationship between Λ -hypernuclei and nuclei...simply because the mass difference of 80 MeV between the Λ and Σ hyperons...is a very large energy relative to the typical energies associated with nuclear excitations.* This difficulty was eliminated by Kerman and Lipkin [28] who suggested in 1971 to consider the Sakata triplet-SU(3) unitary symmetry version in which the proton, neutron and Λ were degenerate. This suggestion was further limited by us in 1976 to $(1p)_{p,n,\Lambda}$ states and, together with Pauli-spin SU(2) symmetry, led to the consideration of Pauli-Sakata SU(6) supermultiplets encompassing nuclei and hypernuclei [29], in direct generalisation of Wigner’s supermultiplet theory of spin-isospin SU(4) symmetry in light nuclei. The analysis of these SU(6) supermultiplets proved very useful for the development of shell model techniques in the 1980s and on by John Millener and collaborators [30]. In particular, the 1976 work focused on the concept of the ‘supersymmetric’ state in addition to the ‘analogue’ state, with the low-lying supersymmetric state arising from the non existence of a Pauli exclusion principle between the Λ hyperon and nucleons.

2.3 Lasting contributions

I wish to highlight two contributions which are likely to remain with us and become textbook chapters in hypernuclear physics.

- (i) Dalitz’s outstanding contribution in the 1960s to weak interactions in hypernuclei, together with Martin Block [31], was to formulate the $\Lambda N \rightarrow NN$ phenomenology of non-mesonic weak-interaction decay modes that dominate the decays of medium-weight and heavy hypernuclei, a process that cannot be studied on free baryons and which offers new systems, Λ hypernuclei, for exploring the little understood $\Delta I = 1/2$ rule in non-leptonic weak interactions. This subject was discussed thoroughly in HYP06 (talks by H. Outa and by G. Garbarino, in these Proceedings) but more experimentation is needed before the underlying physics is fully understood.
- (ii) Another pioneering contribution, in the 1970s, following the introduction of shell-model techniques [32] was to chart the production and γ -ray decay schemes anticipated for excited states in light Λ hypernuclei in order to derive the complete spin dependence of the ΛN interaction effective in these hypernuclei [33]. This work, which I was fortunate to coauthor, was further developed together with John Millener and Carl Dover [34], serving as a useful guide to the hypernuclear γ -ray measurements completed in the last few years, at BNL and at KEK [35], which yielded full determination of the spin dependence in the low-lying spectrum (talks by H. Tamura and by D.J. Millener, in these Proceedings).

3 $\Lambda\Lambda$ hypernuclei

Dalitz in fact anticipated that $\Lambda\Lambda$ hypernuclei be observed and that as a rule they would be particle stable with respect to the strong interaction. His Letter titled *The $\Lambda\Lambda$ -hypernucleus and the $\Lambda - \Lambda$ interaction* [36] appeared as soon as the news of the first observed $\Lambda\Lambda$ -hypernucleus ${}^{10}_{\Lambda\Lambda}\text{Be}$ was reported in 1963 [37] and was followed by a regular paper [38]. He did not work on $\Lambda\Lambda$ hypernuclei for a long period, until 1989, apparently because there were hardly new experimental developments in this field except for the ${}^6_{\Lambda\Lambda}\text{He}$ dubious event reported by Prowse in 1966. He returned to this subject in 1989 [39] feeling the need to scrutinize carefully the interpretation of the ${}^{10}_{\Lambda\Lambda}\text{Be}$ event and its implications in view of a renewed experimental interest to search for the H -dibaryon. This scientific chapter in Dalitz’s life is described in Don Davis’ companion talk in these Proceedings.

4 Σ hypernuclei

Dalitz was puzzled by the CERN-PS low-statistics evidence in the beginning of the 1980, and subsequently by the KEK-PS low-statistics evidence in 1985, for relatively

narrow Σ -hypernuclear peaks in the continuum. The large $\Sigma N \rightarrow \Lambda N$ low-energy cross section, due primarily to the strong pion exchange potential, did not leave much room for narrow Σ states in nuclei; indeed, the first rough estimate by Gal and Dover [40] gave nuclear-matter widths of order $\Gamma_\Sigma \sim 25$ MeV. The suggestion by these authors that some Σ -hypernuclear levels could selectively become fairly narrow due to the $S = 1$, $I = 1/2$ dominance of the $\Sigma N \rightarrow \Lambda N$ transition fascinated him to the extent that he argued favorably for the validity of this interpretation in his 1980 Nature article *Discrete Σ -hypernuclear states* [41], although taking it with a grain of salt. He came back to this subject in 1989, after hearing in HYP88 at Padova Hayano's report of the KEK experiment [42] finding evidence for a $\frac{4}{\Sigma}\text{He}$ near-threshold narrow state. Recalling some old bubble-chamber data on K^- -absorption yields in ^4He near the Σ threshold, he questioned together with Davis and Deloff [43] the compatibility of assigning this $\frac{4}{\Sigma}\text{He}$ as a quasi-bound state with the older data: *Is there a bound $\frac{4}{\Sigma}\text{He}$? He came back to these questions with Deloff in both HYP91 in Shimoda and HYP94 in Vancouver* [44,45].

5 Exotic structures

I have already mentioned that Dalitz was far from jumping on band wagons of speculative ideas unless there were some good experimental or phenomenological tests to be made in a concrete manner. In this context one finds a Nature paper coauthored by Dalitz, *Growing drops of strange matter* [46], discussing a possible scenario for getting into strange quark matter. It is therefore interesting to wonder how Dalitz would have reacted to the flood of recent reports on the possible existence of \bar{K} -nuclear bound states and on the ongoing experimental searches for such objects. The methodology adopted in the KEK and in the Frascati dedicated experiments discussed in the HYP06 conference was to use stopped K^- reactions, partly relying on Akaishi and Yamazaki's production rate estimate of $\sim 2\%$ per stopped K^- in ^4He [47]. This estimate is totally unacceptable since a similar production rate is known to hold at rest for (the most favorable) $A = 4$ hypernuclei [48]; hypernuclei are produced via the dominant absorptive $K^- N \rightarrow \pi Y$ modes, whereas the $K^- N \rightarrow N\bar{K}$ backward-elastic mode responsible for replacing a bound nucleon by a bound \bar{K} is suppressed at rest with respect to the former reactive modes owing to the $1/v$ law near threshold. Realistic estimates should give rates of order 10^{-4} or less, per stopped K^- , for the production of \bar{K} -nuclear bound states. In-flight K^- reactions are more promising, but unfortunately will not be feasible before J-PARC is operated, from 2009 on. Preliminary (K^-, p) and (K^-, n) spectra at $p_{\text{lab}} = 1$ GeV/c on ^{12}C obtained in KEK-E548 show only appreciable strength in the \bar{K} bound-state region, but no peaks [49], in accordance with a recent in-flight reaction calculation [50]. Given this situation, the use of other methods, using proton or antiproton beams, or nucleus-nucleus collisions, has been ad-

vocated. Let me mention briefly some of the recent claims in this rather speculative area.


A preliminary evidence for a broad peak in the Λd invariant-mass spectrum at $M_{\text{inv}}(\Lambda d) = 3159 \pm 20$ MeV, and a width $\Gamma = 100 \pm 50$ MeV, was reported recently by the FOPI detector collaboration at GSI [51] in a study of ΛX correlations ($X = p, d, t, \dots$) in Ni+Ni collisions at 1.93 GeV/A. This is barely compatible with the very narrow peak at 3140 MeV reported in the E471 KEK $^4\text{He}(K^-, n)$ experiment [52] as an evidence for the $I = 0$, $\bar{K}NNN$ deeply bound narrow state predicted by Akaishi and Yamazaki [47] and recently withdrawn (M. Iwasaki, these Proceedings). However, the Λd peak observed in the GSI experiment could be correlated with the Λp relatively narrow peak observed in $\bar{p} - ^4\text{He}$ annihilation at rest by the OBELIX spectrometer collaboration at the LEAR facility in CERN (T. Bressani, these Proceedings and in Ref. [53]) provided it is accompanied by an unseen neutron spectator. It should be noted that the statistical significance of these two peaks that imply deep binding $B_{\bar{K}} \sim 160$ MeV is not particularly high, 4.5 and 4 respectively.²

Recently, the FOPI collaboration at GSI reported a more robust evidence for another peak [54] which naively would be interpreted as due to a deeply bound $K^- pp$, by detecting Λp pairs in both Ni+Ni and Al+Al collisions. Preliminary results are shown in Fig. 1, where the Λp invariant mass peaks near the ΣN threshold, with mass uncertainty of about 20 MeV and with a substantial width. In contrast, the invariant mass corresponding to the signal assigned to $K^- pp$ by the FINUDA spectrometer collaboration [55] is marked by a vertical dashed line across the shown spectra, with no peaking whatsoever. The possibility of a resonance or cusp phenomenon for the Λp system, at or near the opening of the ΣN threshold, has always intrigued Dalitz. However, I dare say that had he been with us today, he would have considered favorably another possibility, that the light, only Σ hypernucleus known to be bound, $\frac{4}{\Sigma}\text{He}$ is the source of these Λp pairs. The binding energy of this hypernucleus is $B = 4.4 \pm 0.3(\text{stat}) \pm 1(\text{syst})$ MeV, with respect to the $\Sigma^+ + ^3\text{H}$ threshold, and the value of width assigned to it is $\Gamma = 7.0 \pm 0.7 + 1.2$ MeV [56]. Its quantum numbers are $I = 1/2$, $J^\pi = 0^+$ [57] with all four baryons in s states. In particular, it may be viewed in isospace as a linear combination of Σ^+ coupled to ^3H and Σ^0 coupled to ^3He . Its wavefunction is schematically given by:

$$\Psi(\frac{4}{\Sigma}\text{He}) = \alpha(\Sigma N)_{I=1/2, 3/2}^{S=0}(NN)_{I=1}^{S=0} + \beta(\Sigma N)_{I=1/2}^{S=1}(NN)_{I=0}^{S=1}, \quad (1)$$

where only the spin-isospin structure is specified. The decay of $\frac{4}{\Sigma}\text{He}$ is dominated by the $(\Sigma N \rightarrow \Lambda N)_{I=1/2}^{S=1}$ two-body transition, proceeding therefore through the component with amplitude β in which the NN composition is pn . This means that the ΣN composition is a mixture of

² Bendiscioli *et al.* [53] also reported a Λd peak with statistical significance ~ 3 at $M_{\text{inv}}(\Lambda d) = 3190 \pm 15$ MeV, with $\Gamma \leq 60$ MeV, which would correspond to an $I = 0$, $\bar{K}NNN$ state bound by about 120 NeV.



Literature:

Strange Dibaryon H_1^+ , $M=2.13$ GeV, $\Gamma=17$ MeV
C. Pigot et al. (Rome-Saclay-Vanderbilt Collaboration), NPB 249 (1985) 172
 Λp resonance in $K^+D \rightarrow \pi^+p\Lambda$ at rest,
 $M_1=2.128$ GeV, $\Gamma_1=7$ MeV and $M_2=2.138$ GeV, $\Gamma_2=9.1$ MeV
Tai Ho Tan, PRL 23, 101 (1969)

Peak position: 2.13 ± 0.02 GeV
Reflection of other resonances?
Work in progress ...

N. Hermann, Uni-HD

24/27

Fig. 1. Λp invariant-mass spectrum taken by the FOPI detector collaboration at GSI in Ni+Ni and Al+Al collisions (figure shown by Paul Kienle at this meeting)

Σ^+n and Σ^0p , both of which decay to Λp . One expects then $\frac{4}{2}\text{He}$ to decay dominantly by emitting back-to-back Λp pairs with slower ‘spectator’ proton and neutron which will somewhat distort the $\Sigma N \rightarrow \Lambda p$ two-body kinematics. A more conclusive proof for this suggestion would come from the observation of back-to-back $\Lambda^3\text{He}$ pairs in the two-body decay $\frac{4}{2}\text{He} \rightarrow \Lambda + {}^3\text{He}$. The branching ratio for this decay relative to the inclusive ΛX decay rate is perhaps a few percent, as may be argued by analogy with the approximately 8%(5%) branching ratio measured for the nonmesonic decay $\frac{4}{\Lambda}\text{He}(\frac{5}{\Lambda}\text{He}) \rightarrow n + {}^3\text{He}(\frac{4}{\Lambda}\text{He})$ relative to the inclusive π^- decay rate of $\frac{4}{\Lambda}\text{He}(\frac{5}{\Lambda}\text{He})$ [58,59]. Irrespective of whether or not the above conjecture of $\frac{4}{2}\text{He}$ production is correct for the FOPI-Detector GSI experiments, it would be a wise practice for \bar{K} -nuclear bound state searches in heavy ion collisions to first look for known hypernuclear signals in order to determine their production rates as calibration and normalization standards.

6 Concluding remarks

Dalitz’s lifelong study of hypernuclei was central to his career as a phenomenologically inclined theoretical physicist. His style was unique. Asked by then his student Chris Llewellyn Smith about ‘new theories’, Dalitz responded

– *My job is not to make theories - it’s to understand the data,*

he saw the theorist’s role as being to find a way of representing experimental data so that they directly reveal nature’s secrets, as the Dalitz Plot had done [60]. His lifelong nourishment of hypernuclei has shaped and outlined for the last 50 years a field that is now maturing into a broader context of Strangeness Nuclear Physics. His wise and critical business-like attitude will be missed as new experimental facilities are inaugurated with the promise of discovering new facets of this field.

References

1. I.J.R. Aitchison, F.E. Close, A. Gal, D.J. Millener, Nucl. Phys. A **771** (2006) 8.
2. M. Danysz, J. Pniewski, Phil. Mag. **44** (1953) 348.
3. R.H. Dalitz, Rev. Mod. Phys. **33** (1961) 471.
4. R.H. Dalitz, Annu. Rev. Nucl. Sci. **13** (1963) 339.
5. R.H. Dalitz, S.F. Tuan, Phys. Rev. Lett. **2** (1959) 425.
6. M. Alston *et al.*, Phys. Rev. Lett. **6** (1961) 698.
7. R.H. Dalitz, T.C. Wong, G. Rajasekaran, Phys. Rev. **153** (1967) 1617.
8. Y. Tomozawa, Nuovo Cimento A **46** (1966) 707.
9. S. Weinberg, Phys. Rev. Lett. **17** (1966) 616.
10. L. Castillejo, R.H. Dalitz, F.J. Dyson, Phys. Rev. **101** (1956) 453.
11. R.H. Dalitz, Strange particle theory in the cosmic-ray period, in *Proc. Colloque Int. sur l’Histoire de la Physique des Particules, Paris 1982*, J. Physique **43**, Colloq. C8 Suppl. (1982) p. 406.

12. M. O'Byrne, From 'tau' to 'top' - the man behind the Dalitz plot, CERN COURIER **44**, issue 2 (2004) article 19.
13. R.H. Dalitz, Nucl. Phys. A **754** (2005) 14c.
14. R.H. Dalitz, Phys. Rev. **99** (1955) 1475.
15. B.W. Downs, R.H. Dalitz, Phys. Rev. **114** (1959) 593.
16. R.H. Dalitz, R.C. Herndon, Y.C. Tang, Nucl. Phys. B **47** (1972) 109.
17. R.H. Dalitz, Nature of the $\Lambda^0 - N$ force from binding energies of light hyperfragments, in *Proc. 6th Annual Rochester Conference on High Energy Nuclear Physics, Rochester 1956*, Eds. J. Ballam *et al.* (Interscience, New York, 1956) pp. V 40-43.
18. R.H. Dalitz, L. Liu, Phys. Rev. **116** (1959) 1312.
19. R. Ammar, R. Levi Setti, W. Slater, S. Limentani, P. Schlein, P. Steinberg, Nuovo Cimento **19** (1961) 20.
20. R. Ammar, W. Dunn, M. Holland, Nuovo Cimento **26** (1962) 840.
21. R.H. Dalitz, R. Levi Setti, Nuovo Cimento **30** (1963) 489.
22. D.H. Davis, J. Sacton, Hypernuclear physics using photographic emulsion, in *Proc. Int. Conf. Hypernuclear Physics, ANL 1969*, Eds. A.R. Bodmer, L.G. Hyman (ANL, Argonne IL, 1969) pp. 159-192.
23. G. Bohm *et al.*, Nucl. Phys. B **24** (1970) 248; M. Juric *et al.*, Nucl. Phys. B **47** (1972) 36.
24. R.H. Dalitz, D.H. Davis, D.N. Tovee, Nucl. Phys. A **450** (1986) 311c.
25. C.B. Dover, A. Gal, G.E. Walker, R.H. Dalitz, Phys. Lett. B **89** (1979) 26.
26. R.H. Dalitz, A. Gal, Phys. Lett. B **64** (1976) 154.
27. R.H. Dalitz, The present problems and future outlook in Λ -hypernuclear physics, in *Proc. Int. Conf. Hypernuclear Physics, ANL 1969*, Eds. A.R. Bodmer, L.G. Hyman (ANL, Argonne IL, 1969) pp. 708-747.
28. A.K. Kerman, H.J. Lipkin, Ann. Phys. **66** (1971) 738.
29. R.H. Dalitz, A. Gal, Phys. Rev. Lett. **36** (1976) 362.
30. E.H. Auerbach, A.J. Baltz, C.B. Dover, A. Gal, S.H. Kahana, L. Ludeking, D.J. Millener, Phys. Rev. Lett. **47** (1981) 1110; Ann. Phys. **148** (1983) 381.
31. M.M. Block, R.H. Dalitz, Phys. Rev. Lett. **11** (1963) 96.
32. A. Gal, J.M. Soper, R.H. Dalitz, Ann. Phys. **63** (1971) 53; **72** (1972) 445; **113** (1978) 79.
33. R.H. Dalitz, A. Gal, Ann. Phys. **116** (1978) 167.
34. D.J. Millener, A. Gal, C.B. Dover, R.H. Dalitz, Phys. Rev. C **31** (1985) 499.
35. O. Hashimoto, H. Tamura, Prog. Part. Nucl. Phys. **57** (2006) 564.
36. R.H. Dalitz, Phys. Lett. **5** (1963) 53.
37. M. Danysz *et al.*, Phys. Rev. Lett. **11** (1963) 29; Nucl. Phys. **49** (1963) 121.
38. R.H. Dalitz, G. Rajasekaran, Nucl. Phys. **50** (1964) 450.
39. R.H. Dalitz, D.H. Davis, P.H. Fowler, A. Montwill, J. Pniewski, J.A. Zakrzewski, Proc. Royal Soc. A **426** (1989) 1.
40. A. Gal, C.B. Dover, Phys. Rev. Lett. **44** (1980) 379.
41. R.H. Dalitz, Nature **285** (1980) 11.
42. R.S. Hayano *et al.*, Nuovo Cimento A **102** (1989) 437.
43. R.H. Dalitz, D.H. Davis, A. Deloff, Phys. Lett. B **236** (1990) 76.
44. R.H. Dalitz, A. Deloff, Nucl. Phys. A **547** (1992) 181c.
45. R.H. Dalitz, A. Deloff, Nucl. Phys. A **585** (1995) 303c.
46. G.L. Shaw, M. Shin, R.H. Dalitz, M. Desai, Nature **337** (1989) 436.
47. Y. Akaishi, T. Yamazaki, Phys. Rev. C **65** (2002) 044005.
48. D.H. Davis, Nucl. Phys. A **754** (2005) 3c.
49. T. Kishimoto, private communication (2006).
50. J. Yamagata, H. Nagahiro, S. Hirenzaki, Phys. Rev. C **74** (2006) 014604.
51. N. Herrmann, invited talk presented at EXA 2005, Vienna (<http://www.oeaw.ac.at/smi/exa05/program.htm>).
52. T. Suzuki *et al.*, Nucl. Phys. A **754** (2005) 375c.
53. G. Bendiscioli, A. Fontana, L. Lavezzi, A. Panzarasa, A. Rotondi, T. Bressani, submitted for publication in Nucl. Phys. A (2006).
54. P. Kienle, private communication (2006).
55. M. Agnello *et al.*, Phys. Rev. Lett. **94** (2005) 212303.
56. T. Nagae *et al.*, Phys. Rev. Lett. **80** (1998) 1605.
57. T. Harada, Phys. Rev. Lett. **81** (1998) 5287.
58. G. Coremans *et al.*, Nucl. Phys. B **16** (1970) 209.
59. G. Keyes, J. Sacton, J.H. Wickens, M.M. Block, Nuovo Cimento A **31** (1976) 401.
60. C.H. Llewellyn Smith, remarks at the Dalitz memorial meeting at the University of Oxford, 3 June 2006.